



Extension FactSheet

Department of Entomology, 1991 Kenny Road, Columbus, Ohio 43210

Carpenter Bees

Susan C. Jones

Associate Professor, Entomology; Extension Specialist, Household and Structural Pests

Carpenter bees are so named because they excavate galleries in wood to create nest sites. They do not consume wood. Rather, they feed on pollen and nectar. Carpenter bees are important pollinators of flowers and trees. Carpenter bees typically are just nuisance pests that cause cosmetic rather than structural damage to wood. Nonetheless, considerable wood damage can result from many generations of carpenter bees enlarging existing galleries in wood.

Large carpenter bees belong to the genus *Xylocopa*. Two native species, *Xylocopa virginica* and *Xylocopa micans*, occur in the eastern United States. There also are a number of native carpenter bees in the western United States. This fact sheet primarily pertains to *X. virginica*, which has the common name of carpenter bee.

Identification

Carpenter bees are large and robust. *X. virginica* is three-fourths to one-inch long, black, with a metallic sheen. The thorax is covered with bright yellow, orange, or white hairs, and the upper side of the abdomen is black, glossy, and bare (Figure 1). The female has a black head, and the male has white markings on the head. Carpenter bees have a dense brush of hairs on the hind legs.

Carpenter bees somewhat resemble bumble bees, except bumble bees have dense yellow hairs on the abdomen and large pollen baskets on the hind legs. Various species of bumble bees and carpenter bees are similar in size. Bumble bees typically nest in the ground whereas carpenter bees nest in wood.

Life History

Carpenter bees are solitary insects that do not form colonies. Male and female carpenter bees overwinter as adults within their old nest gallery. Adults emerge in



Figure 1. Carpenter bee.
(Courtesy of Kansas State University.)

the spring (April and early May) and mate. There is one generation per year.

The males are not long lived, and the female carpenter bee prepares the nest. Gallery construction is a time- and energy-consuming process, and the female will preferentially refurbish an old nest rather than excavate a new one. When constructing a new nest, the female uses her strong jaws (mandibles) to excavate a clean-cut, round nest entrance hole on the lateral surface of wood in an exposed or unexposed location. This hole is slightly less than 1/2-inch wide, approximately the diameter of her body. She bores into the wood perpendicular to the grain for one to two inches then makes a right angle turn (~90°) and excavates along the wood grain for four to six inches to create a gallery (tunnel). She excavates the gallery at the rate of about one inch in six days.

The female bee creates a series of provisioned brood cells in the excavated gallery. The larval provision consists of a mixture of pollen and regurgitated nectar formed into a ball. The female forms a food ball at the far end of an excavated gallery, lays an egg on top of the mass, and then walls off the brood cell with a plug of chewed wood pulp. A female often creates six to 10 partitioned brood cells

in a linear row in one gallery, and she dies soon thereafter. Larvae feed on the pollen/nectar food mass, which is sufficient food for them to develop to the adult stage.

The life cycle (egg, larva, pupa, adult) is completed in approximately seven weeks, although developmental time may vary depending on the temperature. The new adults typically remain in their gallery for several weeks then chew through the cell partitions and venture outside in late August. They collect and store pollen in the existing galleries, but also spend much of their time just huddled together inside a gallery. These new adults hibernate in galleries because they require shelter during the winter. They then emerge the following spring.

Habits

Carpenter bees nest in a wide range of softwoods and hardwoods, particularly if the wood is weathered. Eastern species of carpenter bees prefer softwoods such as cedar, redwood, cypress, pine, and fir. The bees can more easily tunnel through woods that are soft and that have a straight grain. Western species of carpenter bees often nest in oak, eucalyptus, and redwood.

Carpenter bees attack structural timbers and other wood products, including fence posts, utility poles, firewood, arbors, and lawn furniture. In buildings, carpenter bees nest in bare wood near roof eaves and gables, fascia boards, porch ceilings, decks, railings, siding, shingles, shutters, and other weathered wood. These bees avoid wood that is well painted or covered with bark.

The carpenter bee entrance hole in wood may not necessarily be in an exposed area. For example, the inner lip of fascia boards is a common site of attack. Nail holes, exposed saw cuts, and unpainted wood are attractive sites for the bees to start their excavations.

Economic Importance

Despite their beneficial aspect of being important pollinators of many trees and flowers, carpenter bees also may be nuisance pests around structures. Carpenter bees are noisy, which may be bothersome. These large bees create alarm when they dive-bomb or fly erratically around humans. In actuality, these are male bees, which are territorial but harmless because they lack a stinger. Only females have a stinger. Female carpenter bees are docile and are reported to sting only if handled.

Carpenter bees create a nuisance by excavating round entry holes in wood (Figure 2) and depositing yellowish to brownish streaks of excrement and pollen on surfaces below entry holes (Figure 3). They also produce coarse

sawdust from their borings. The carpenter bee gallery system is confined within the wood (Figure 4) and hence is not visible.

Carpenter bee damage to wood initially is minor, and carpenter bees seldom cause consequential structural damage. However, their repeated colonization of the same wood can eventually cause considerable wood damage. Carpenter bees preferentially refurbish and enlarge an existing tunnel instead of boring a new one, and a gallery can extend for 10 feet if used by many carpenter bees over the years.



Figure 2. Carpenter bee entrance hole in fascia.



Figure 3. Carpenter bee staining on siding below the fascia.



Figure 4. Carpenter bee gallery exposed in wood.

Carpenter bees sometimes construct new tunnels near old ones, with infestations persisting for several years. This complex system of tunnels can result in extensive damage to wood. Wood replacement is necessary when the strength of structural members, posts, poles, and other wood products is reduced due to carpenter bee damage.

Carpenter bees also may be indirectly responsible for unsightly wood damage when woodpeckers riddle the wood with holes searching for the developing carpenter bees to feed upon.

Integrated Pest Management

When dealing with carpenter bees, it is preferable to locate tunnel entrances during the daytime, but treat after dark on a cool evening when carpenter bees are less active. Wear protective clothing to avoid any stings during treatment.

Prevention

Keep all exposed wood surfaces well painted with a polyurethane or oil-base paint to deter attack by carpenter bees. Periodically inspect painted surfaces, because the coatings will begin to deteriorate due to weathering, leaving exposed wood that the bees then can easily attack. Wood stains will not prevent damage. Consider using aluminum, asbestos, asphalt, vinyl siding, and similar non-wood materials that are not damaged by carpenter bees. Seal existing gallery entrance holes to discourage carpenter bees that are looking for potential nesting sites.

Mechanical Measures

A non-insecticidal management approach is to deny carpenter bees access to their galleries by sealing each entrance hole. Thoroughly plug the hole with caulking compound, wood putty, or a wooden dowel affixed with wood glue. If possible, also fill the entire gallery system with a sealant. Carpenter bee galleries are a critical resource, since the bees spend much of their time inside a gallery, and they require its protective conditions to survive the winter. Bees that are trapped inside a caulked gallery typi-

cally will not chew out due to behavioral constraints. This barrier approach has promise for reducing future carpenter bee infestations.

In new nests, the single female often can be swatted and killed, or she can be captured and crushed or otherwise destroyed. Larvae and pupae can be killed by inserting a sturdy wire into the entrance hole and probing into the gallery as deeply as possible.

Insecticides

A chemical treatment using an appropriately labeled insecticide can protect wood for short periods, especially in the spring and summer when carpenter bee nesting activity is apparent. Dust formulations typically provide residual effects and are effective due to the nature of carpenter bee gallery construction. Precisely inject the dust directly into each nest entrance hole and as deep into the tunnel as possible and also apply it to the adjacent wood surface. Wait for a few days before plugging entrance holes since adult bees should be allowed to pass freely to distribute the insecticide within the galleries. Newly emerged bees also will contact the dust when attempting to leave their gallery.

For use as a preventive, an insecticide should be applied to wood in early spring before carpenter bees begin excavating nests. The insecticide kills the bees that contact it on the wood's surface. However, a preventive approach has limitations because of the difficulty in applying a chemical to all exposed wood on the house where bees could nest. Furthermore, such insecticides usually degrade in a matter of weeks or months so repeated applications are needed to maintain a lethal dose of the insecticide. Some pest management companies report good results against carpenter bees by spraying wood with a microencapsulated pyrethroid, Demand® CS insecticide (registered for use only by licensed professional applicators), which contains the active ingredient lambda-cyhalothrin. A number of other pyrethroids (bifenthrin, cyfluthrin, deltamethrin, permethrin, etc.) also are labeled for use against carpenter bees.

Insecticides that act as stomach poisons, such as borates, typically are ineffective against carpenter bees, which do not ingest the wood that they excavate.

Visit Ohio State University Extension's web site "Ohioline" at <http://ohioline.osu.edu>

OSU Extension embraces human diversity and is committed to ensuring that all educational programs conducted by Ohio State University Extension are available to clientele on a nondiscriminatory basis without regard to race, color, age, gender identity or expression, disability, religion, sexual orientation, national origin, or veteran status.

Keith L. Smith, Associate Vice President for Agricultural Administration and Director, OSU Extension
Copyright © 2006, The Ohio State University